

Tips for Managing Complex Regional Pain Syndrome

September 11, 2015 by [Jim Ducharme, MD, CM, FRCPC](#)



Hands of a patient with early CRPS, before atrophic changes set in.

Complex regional pain syndrome (CRPS), previously known as reflex sympathetic dystrophy (RSD), is a chronic neuropathic pain condition that can arise from trauma of any kind. It can be the result of something as minor as a blood draw that initiates a reaction. The condition arises more frequently than many emergency physicians may realize: roughly 3 percent of patients suffering a Colles fracture develop CRPS. Often the traumatic event cannot be remembered, and CRPS has been associated with trauma happening anywhere from a day to a year after the event. Its most consistent feature, however,

is how often physicians fail to make the diagnosis on initial presentation. Furthermore, our lack of understanding about how to manage the severe pain that occurs during acute flare-ups of this chronic condition worsens the suffering that many patients with CRPS endure over decades.

Early Diagnosis Is Key

As can be seen with diabetic neuropathy, CRPS has both a sensory and an autonomic dysfunction. Unlike patients with diabetic neuropathy, both will be present from the start. Not only do patients suffer from intense pain that does not correspond to a specific nerve distribution, they also suffer visible changes as the result of their autonomic dysfunction.

Initially, the involved painful area (usually part of an extremity) becomes red, warm, and edematous; it is often initially misdiagnosed as cellulitis. The presence of severe allodynia (pain induced with a nonpainful stimulus such as light touch) should make the physician consider the true diagnosis. It is very important that CRPS be diagnosed early on because active treatment can reverse and eliminate the condition. Treatment includes neuropathic analgesics (eg, tricyclics, gabapentinoids) combined with active physiotherapy and mindfulness. Many patients who develop this condition will come to the emergency department with their painful condition when it begins, so the emergency physician needs to be able to diagnose and refer appropriately. I personally diagnose two to three new cases per year in my emergency medicine practice.



Feet of a patient with more advanced CRPS, which show clear skin atrophic changes.

Failure to treat within the first weeks of symptom onset will allow the physical changes to start. The involved area will develop dystrophic skin changes: a shiny, thin, erythematous appearance. Underlying muscles atrophy so that the involved area becomes wasted in appearance over time. Typical burning neuropathic pain persists. If left untreated (or if poorly treated), CRPS can spread, involving larger parts of the body.

Treating Flare-ups

Patients will also present to the emergency department because of an acute flare-up of their chronic pain. CRPS can become acutely more painful because of N-methyl-D-aspartate (NMDA) activity and hyperresponsiveness to NMDA. NMDA is a neurotransmitter present in the dorsal horns and spinothalamic tracts, and it is the number-one initiator of wind-up in acutely painful conditions. With CRPS flare-ups, it is almost as if wind-up starts over again. The burning pain becomes acutely worse; pain is severe and unresponsive to almost all analgesics. Opioids will not control the pain of a flare-up unless given in a quantity that would make the patient somnolent. Opioids should not be considered a first-line treatment in this situation. It is recognized that many patients with CRPS ask for opioids for their severe pain. As with any patient asking for opioids when suffering from a chronic pain condition, this can create distrust and a stressful environment. Increasingly, national patient groups are educating patients that opioids will not be effective.

Given the cause of the pain flare-up, the treatment needs to be directed at stopping the NMDA activity. This is best accomplished with ketamine, an NMDA antagonist. A patient can only receive intravenous ketamine in a hospital environment, so emergency physicians need to be able to recognize and treat these severe pain flare-ups.

Treatment Is Straightforward:

1. Initial bolus of 0.2–0.3 mg/kg of ketamine infused over 10 minutes. Giving this dose as an IV push will produce a high rate of dissociative side effects (up to 75 percent of patients) and should be avoided. Almost diagnostic is the patient's response: severe pain should be resolved by the end of the 10-minute bolus.
2. An infusion of ketamine (0.2 mg/kg/hr) for four to six hours. Although the medical literature for this is almost nonexistent, clinical experience has shown that an infusion of this duration resets the NMDA activity to baseline. Patients can return home on their usual medications, with the expectation that the flare-up, which can normally last weeks, will be over. Return rates for the same flare-up after ketamine treatment approach zero. For readers who feel four to six hours is too long, I encourage them to try shorter periods (two or three hours) and publish their results. No discharge prescription from the emergency department will be required.

Patients do not require admission, and they should not receive opioids. They do require the acute ketamine intervention, or they will suffer severe pain for weeks as a result of the flare-up. To date, there is no other effective treatment for a CRPS pain flare-up. Some researchers have studied an infusion of 5 mg/kg of lidocaine over a 60-minute period as an alternative treatment plan, but results are variable. Referral of newly diagnosed patients to physiotherapy and a comprehensive pain program is critical.

With better understanding of CRPS, emergency physicians will know when and how to intervene. Concern over drug seeking should be allayed, allowing appropriate care to be provided.